

Measure Descriptions

This table reflects all measures included in CHPI reporting. Measures reported for individual physicians are a subset of these. In the Data Type field, a “C” indicates that the measure includes Commercial data, and an “M” indicates that the measure includes Medicare data.

NQF #	Measure Name	Measure Description	Steward	Data Type
N/A	Diabetes Care Composite	The simple average of an individual physician’s or practice site’s scores on: Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Testing (NQF #0057), Comprehensive Diabetes Care: Medical Attention for Nephropathy (NQF #0062), Diabetes Care: Appropriate Treatment of Hypertension (NQF #0546)	N/A	C,M
0002	Appropriate Testing for Children with Pharyngitis	The percentage of children 2–18 years of age who were diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus (strep) test for the episode	NCQA	C
0057	Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Testing	Percentage of patients aged 18 through 75 years of age with diabetes who had hemoglobin A1c (HbA1c) testing	NCQA	C,M
0062	Comprehensive Diabetes Care: Medical Attention for Nephropathy	The percentage of members 18-75 years of age with diabetes (type 1 and/or type 2) who received a nephropathy screening test or had evidence of nephropathy	NCQA	C,M
0069	Appropriate Treatment for Children with Upper Respiratory Infection	Percentage of children 3 months to 18 years of age with a diagnosis of URI who were not dispensed an antibiotic medication. The measure is reported as an inverted rate [1 – (numerator/eligible population)]. A higher score indicates appropriate treatment of upper respiratory infection (i.e., the proportion who were not prescribed or dispensed an antibiotic prescription).	NCQA	C,M
0541	Proportion of Days Covered (PDC)	The percentage of patients 18 years and older who met the proportion of days covered (PDC) threshold of 80%. A performance rate is calculated separately for the following medication categories: Renin Angiotensin System (RAS) Antagonists, Diabetes Medications, Statins.	PQA	C,M
0546	Diabetes Care: Appropriate Treatment of Hypertension	The percentage of patients who were dispensed a medication for diabetes and hypertension that are receiving an angiotensin-converting -enzyme-inhibitor (ACEI) or angiotensin receptor blocker (ARB) or direct renin inhibitor (DRI) renin-angiotensin-antagonist medication	PQA	C,M
1516	Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life	The percentage of members 3–6 years of age who had one or more well-child visits with a PCP	NCQA	C
N/A	Adolescent Well-Care Visits	The percentage of enrolled members 12–21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner	NCQA	C
N/A	Breast Cancer Screening – Ages 50-69	The percentage of women ages 50 to 69 who had one or more mammograms	NCQA	C,M
N/A	Annual Monitoring for Patients on Persistent Medications_Total_ACE inhibitors or ARBs or Diuretics	The percentage of members 18 years of age and older who received at least 180 treatment days of ambulatory medication therapy and at least one therapeutic monitoring event for the therapeutic agent. Report each of the three rates separately and as a total rate. <ul style="list-style-type: none"> • Annual monitoring for patients on angiotensin converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARB) • Annual monitoring for patients on diuretics 	NCQA	C,M